arXiv:1207.4623v1 [cond-mat.stat-mech] 19 Jul 2012

Statistics of entanglement production in a chaotic dot with non-ideal contacts

Dario Villamaina and Pierpaolo Vivo
Laboratoire de Physique Théorique et Modèles Statistiques (UMR 8626 du CNRS),
Université Paris-Sud, Bâtiment 100, 91405 Orsay Cedex (France)
(Dated: July 20, 2012)

We compute analytically the full distribution of concurrence for the production of electronic entanglement in a chaotic quantum dot. The dot is connected to the external world via two leads, one of which is assumed to be ideal while the other is partially transparent. While previous works could address the issue only by means of numerical simulations, our analytical predictions stem from the recent calculation [P. Vidal and E. Kanzieper, *Phys. Rev. Lett.* **108**, 206806 (2012)] of the joint distribution of transmission eigenvalues in non-ideal cavities for the case of broken time-reversal symmetry. The analytical results are valid for any value of the transparency parameter of the non-ideal lead. They are in excellent agreement with numerical simulations and imply that highly asymmetric transparencies for the contacts in the two leads significantly enhance the probability of producing entangled states inside the cavity.

PACS numbers: 03.67.Bg,73.63.Kv,73.23.-b,05.60.Gg
Keywords: Chaotic cavities, quantum dots, entanglement, concurrence

The phenomenon of entanglement between spatially separated particles is probably one of the most baffling predictions of quantum mechanics. Initially, the idea that entangled quantum states could be subject to non-classical correlations was seen as a possible evidence of the incompleteness of quantum theory [1]. Nowadays, thanks to fascinating experimental developments, the reality of quantum entanglement is no longer disputed. Besides its fundamental and intellectually challenging interest, modern research on entanglement mainly focuses on its possible applications to quantum information processing, cryptography and teleportation [2, 3]. As a consequence, it is of paramount interest to discover feasible pathways for the production, manipulation, and detection of quantum entanglement in a variety of physical devices.

Among the most promising routes, the entanglement in solid-state electronic systems [4, 5] plays a prominent role. As far as the production of entangled electrons is concerned, several proposals based on interacting [6] and noninteracting [7–9] electron mechanisms are already available. A very interesting interactionless mechanism was recently proposed by Beenakker *et al.* [9] where a ballistic quantum dot (see sketch in Fig. 1) is used as an orbital entangler for pairs of noninteracting electrons [10]. A quantum dot is basically a mesoscopic electronic billiard connected to the external world by two double-channel leads and brought out of equilibrium by an applied external voltage. Assuming that the classical electron dynamics inside the cavity is chaotic, experimental observables such as conductance and shot noise fluctuate from sample to sample and their statistics (about which nearly everything is known [11]) is governed by the transmission eigenvalues of the cavity.

In order to quantify the degree of entanglement produced inside the cavity, one uses a standard measure called *concurrence*, which is related to the violation of a Bell inequality for current correlators. The average and variance of the concurrence for a chaotic dot *with ideal leads* (i.e. when the tunnelling probability across the leads is unity) was computed in [9]. It was found that these two moments are practically unaffected by the breaking of time-reversal invariance (TRI), but the fluctuations can be very large (of the same order as the average). This implies that a statistical description of entanglement production based on just the first moments is highly inaccurate. The full distribution of the concurrence (again in the case of ideal leads) was computed by Gopar and Frustaglia [12], and indeed they found remarkable differences in the distributions between the cases with preserved $(\beta = 1)$ and broken $(\beta = 2)$ TRI (here β denotes the Dyson index of the cavity), although the first moments are only very mildly affected. Still in the domain of ideal cavities, geometrical constraints on the entanglement production were discovered in [13] implying that more entangled states are less likely to be detected.

All available analytical results heavily rely on the assumption that all leads are ideal. This restriction is however hardly tolerable, as ideal transparencies are never realized in experiments. The tunnelling probabilities (Γ_ν, $\nu = 1, 2$) between the leads and the cavity can be tuned, and it is therefore of great theoretical interest to study the entanglement production as a function of Γ_ν. The available results for non-ideal dots, however, are only numerical and limited to ι) the distribution of charge cumulants [14] and $\iota\iota$) the concurrence distribution [15] for the cases of preserved and broken TRI. The latter investigation led to the interesting outcome that extremely uneven contact transparencies may confer advantages in the production of entanglement, especially for $\beta = 2$. An analytical approach to this problem has been so far hindered by the lack of an explicit formula for the joint distribution of transmission eigenvalues in the case of non-ideal leads. Such an *impasse*, however, can now be overcome thanks to a recent breakthrough by Vidal and Kanzieper [16], who were able to compute the desired joint distribution for the case of a $\beta = 2$ cavity with n_L non-ideal channels in the left lead and n_R ideal channels in the right lead. Using this recent result as a starting point, in this letter we present the first exact analytical calculation for the full distribution and arbitrary moments of the concurrence for $\beta = 2$ (broken TRI) in cavities supporting one ideal and one non-ideal lead, therefore giving a solid support to the previous numerical investigations [15]. The known results in the ideal setting are easily recovered as a limiting case of our theory. Our central result for the concurrence distribution in Eq. (8), expressed in a very compact form, allows to conclude that highly asymmetric transparencies for the contacts in the two leads significantly enhance the probability of producing entangled states with respect to the ideal case. We believe that a confirmation of this prediction may be within reach of current experimental capabilities.

The orbital entangler, first proposed in Ref. [9] is sketched in Fig. 1. It consists of a quantum dot with two attached double-channel leads at the left and right. Each lead is connected to an electron reservoir. Applying a small voltage between the two reservoirs gives rise to a electronic current flowing through the dot from left to right. The scattering within the dot leads to the production of entangled pairs between transmitted (to the right) and reflected (to the left) electrons. More precisely, the outgoing state $|\psi_{\mathrm{out}}\rangle$ of two scattered electrons can be written as the superposition $|\psi_{\mathrm{out}}\rangle = |\psi_{\ell\ell}\rangle + |\psi_{rr}\rangle + |\psi_{\ell r}\rangle$, where $|\psi_{\ell\ell}\rangle$ and $|\psi_{rr}\rangle$ are separable states, corresponding to both electrons being scattered to the left or to the right, while $|\psi_{\ell r}\rangle$ (representing a state where one electron is scattered to the left and the other to the right) may be nonseparable. Since an electron leaving the quantum dot to the left side can choose between channels 1 and 2 for escaping, this defines a two-level quantum system or *qubit*. The same happens with an electron escaping to the right side through leads 3 and 4. This means that the state $|\psi_{\ell r}\rangle$ in general describes (up to a normalization factor) a two-qubit entangled state.



FIG. 1: Sketch of the orbital entangler.

A widely used measure for quantifying two-qubit entanglement is the concurrence $\mathcal{C}$ [17], an entanglement monotone constructed from the two-qubit density matrix. The concurrence is intimately connected to the transport properties of the cavity, and the crucial question is then how to compute it in the proposed setting.

Consider the 4×4 scattering matrix $\mathcal{S}$ of the cavity,

$$\mathcal{S} = \begin{bmatrix} \mathbf{r} & \mathbf{t}' \\ \mathbf{t} & \mathbf{r}' \end{bmatrix} , \tag{1}$$

where $\mathbf{r}, \mathbf{r}', \mathbf{t}$, and $\mathbf{t}'$ are 2×2 reflection and transmission matrices, respectively. In the presence of TRI, $\mathcal{S}$ is unitary and symmetric. If TRI is broken (due to, e.g., the application of a magnetic flux), then $\mathcal{S}$ is only unitary. Let τ_1 and τ_2 be the eigenvalues ($0 \leq \tau_i \leq 1$) of the Hermitian matrix $\mathbf{t}\mathbf{t}^\dagger$. Then, the concurrence $\mathcal{C}$ can be written in terms of the transmission eigenvalues τ_1 and τ_2 as [4, 9]

$$\mathcal{C} = \frac{2\sqrt{\tau_1(1-\tau_1)\tau_2(1-\tau_2)}}{\tau_1 + \tau_2 - 2\tau_1\tau_2}. \tag{2}$$

We note that the concurrence varies from 0 to 1. The case $\mathcal{C} = 0$ corresponds to separable nonentangled states, while maximally entangled (Bell) states correspond to $\mathcal{C} = 1$. Those states with a $0 < \mathcal{C} < 1$ are non-separable partly entangled states. From Eq. (2), the entanglement is maximal ($\mathcal{C} = 1$) when $\tau_1 = \tau_2$, and minimal ($\mathcal{C} = 0$) when $\tau_1 = 0$ and $\tau_2 = 1$ or $\tau_1 = 1$ and $\tau_2 = 0$. A finite value of $\mathcal{C}$ guarantees that the left and right outgoing channels are orbitally entangled.

The chaotic scattering in the cavity implies that the entanglement production is basically a stochastic process, governed by a random scattering matrix. In particular, the concurrence and any other entanglement quantifiers are random variables whose statistics is the central object of this Letter. What is therefore the probability measure to impose on $\mathcal{S}$? It is well established that:

- in the case of *ideal* leads, the distribution of $\mathcal{S}$ is *uniform* within the unitary group: in the presence of TRI, the statistics of an ensemble of unitary and symmetric $\mathcal{S}$ matrices is described by the so-called Circular Orthogonal Ensemble ($\beta = 1$). When TRI is absent, the statistical properties of $\mathcal{S}$ are described by the Circular Unitary Ensemble ($\beta = 2$). The uniformity of $\mathcal{S}$ induces a non-trivial joint density of the transmission eigenvalues $\tau_{1,2}$ of the transmission matrix $\mathbf{t}\mathbf{t}^\dagger$ [18, 19], given by the Jacobi ensemble of random matrices:

  $$P_{0,0}(\tau_1, \tau_2) = c_\beta |\tau_1 - \tau_2|^\beta (\tau_1\tau_2)^{\beta/2-1}, \tag{3}$$

  where $c_1 = 3/4$ and $c_2 = 6$ are normalization constants.

- in the case of *non-ideal* leads supporting n_{L} (left) and n_{R} (right) propagating channels, the distribution of $\mathcal{S}$ is instead given by the so-called Poisson kernel [20],

  $$P_\beta(\mathcal{S}) \propto \left[\det(\mathbb{1}_N - \bar{\mathcal{S}}\mathcal{S}^\dagger)\det(\mathbb{1}_N - \mathcal{S}\bar{\mathcal{S}}^\dagger)\right]^{\beta/2-1-\beta N/2}. \tag{4}$$

  The $N = n_{\mathrm{L}} + n_{\mathrm{R}}$ eigenvalues $\hat{\gamma} = \mathrm{diag}(\{\sqrt{1-\Gamma_j}\})$ of $\bar{\mathcal{S}}$ characterize couplings between the cavity and the leads in terms of tunnel probabilities Γ_j of the j-th electron channel in the leads. In the ideal limit $\Gamma_i \to 1$, we recover uniformity of $\mathcal{S}$ within the unitary group. In contrast with the ideal case, however, no general derivation exists of the joint density of transmission eigenvalues in the non-ideal case. However, for $\beta = 2$ in the special case where the left lead supports n_{L} propagating modes characterized by a set of tunnel probabilities $\hat{\mathbf{\Gamma}}_{\mathrm{L}} = (\Gamma_1, \ldots, \Gamma_{n_{\mathrm{L}}})$, while the right lead (supporting $n_{\mathrm{R}} \geq n_{\mathrm{L}}$ open channels) is kept ideal so that $\hat{\mathbf{\Gamma}}_{\mathrm{R}} = (\Gamma_{n_{\mathrm{L}}+1}, \ldots, \Gamma_{n_{\mathrm{R}}}) = \mathbb{1}_{n_{\mathrm{R}}-n_{\mathrm{L}}}$, the joint probability density function $P_{(\hat{\gamma}_{\mathrm{L}}|\,\mathbf{0})}(\boldsymbol{R})$ of reflection eigenvalues $\{R_j = 1 - \tau_j\}$ equals [16]

$$P_{(\hat{\gamma}_{\rm L}\,|\,\mathbf{0})}(R_1,\ldots,R_{n_{\rm L}}) \propto \prod_{j<k}(R_j-R_k)\det\Big[{}_2F_1(n_{\rm R}+1,n_{\rm R}+1;1;\,\gamma_j^2R_k)\Big]_{(j,k)\in(1,n_{\rm L})}\prod_{j=1}^{n_{\rm L}}(1-R_j)^{\nu}. \tag{5}$$

where $\nu = |n_{\rm L}-n_{\rm R}|$ and $\{\gamma_1^2 = 1-\Gamma_1,\ldots,\gamma_{n_{\rm L}}^2 = 1-\Gamma_{n_{\rm L}}\}$ is a set of $n_{\rm L}$ coupling parameters related to the associated tunnel probabilities in the non-ideal leads, whilst ${}_pF_q$ is the Gauss hypergeometric function.

In order to derive the distribution of the concurrence, it is necessary to specialize Eq. (5) to the case of two transmission eigenvalues ($n_{\rm L} = n_{\rm R} = 2$) and for $\gamma_1 = \gamma_2 = \gamma$, since both channels in the left lead clearly need to have the same transparency. By carefully taking this limit we get

$$P_{\gamma,\gamma}(\tau_1,\tau_2) = \frac{(\gamma^2-1)^8\sum_{i,j=0}^{4}A_{ij}(\gamma)(1-\tau_1)^i(1-\tau_2)^j}{\left(1-\gamma^2(1-\tau_1)\right)^6\left(1-\gamma^2(1-\tau_2)\right)^6} \tag{6}$$

where the matrix $A_{ij}(\gamma)$ is reported in [21].

The distribution of the concurrence is given by definition by:

$$P(\mathfrak{C}) = \left\langle \delta\left(\mathfrak{C} - \frac{2\sqrt{\tau_1(1-\tau_1)\tau_2(1-\tau_2)}}{\tau_1+\tau_2-2\tau_1\tau_2}\right)\right\rangle \tag{7}$$

where the average is taken with respect to the measures (3) or (6). Evaluating the double integral, we eventually get to the surprisingly simple result:

$$P(\mathfrak{C}) = \sum_{k=0,4,6,8} c_k(\mathfrak{C})\gamma^k \tag{8}$$

where the coefficients $c_k(\mathfrak{C})$, expressed only in terms of elementary functions, are listed explicitly in [21].

The limit $\gamma \to 0$, corresponding to ideal cavities, yields

$$P(\mathfrak{C}) = c_0(\mathfrak{C}), \tag{9}$$

which is exactly the probability of concurrence computed in [12], Eq. (15). Let us note that the simple polynomial structure of Eq. (8), not evident in Eq. (6), is essentially due to some symmetry of the integrand. In order to clarify this point, it is useful to write down the expansion in γ of Eq. (6)

$$P_{\gamma,\gamma}(\tau_1,\tau_2) = \sum_{k=0}^{+\infty}\gamma^{2k}P^{(2k)}(\tau_1,\tau_2). \tag{10}$$

Then, clearly $P^{(0)}(\tau_1,\tau_2)$ is given by Eq. (6) and all the odd terms vanish. $P^{(2)}(\tau_1,\tau_2)$ is antisymmetric under the transformation $\tau_1 \to 1-\tau_1$ and $\tau_2 \to 1-\tau_2$, while the concurrence is symmetric. This symmetry explains why all the moments of $\mathfrak{C}$ vanish up to the second order in γ and the first correction in Eq. (8) is $\mathcal{O}(\gamma^4)$. On the contrary, there is no trivial symmetry that kills all higher order terms and this enormous simplification can be explained only by a non-linear change of variables [21]. Moments of any order can be analytically computed from Eq. (8) and the first two are plotted in Fig. 3.

In order to check the analytical predictions, we use the same simple numerical algorithm described in detail in Ref. [15]. The 4×4 scattering matrix of the non-ideal system $\mathcal{S}$ can be decomposed as $\mathcal{S} = \mathcal{R} + \mathcal{T}(\mathbf{1}-\mathcal{S}_0\mathcal{R})^{-1}\mathcal{S}_0\mathcal{T}$ where $\mathcal{S}_0$ is a random matrix distributed uniformly within the unitary group and represents the scattering matrix of an ideal cavity. The matrices $\mathcal{R}$ and $\mathcal{T}$ include information on the transparency of the contacts, as follows:

$$\mathcal{R} = \mathrm{diag}(\mathrm{i}\sqrt{1-\gamma},\mathrm{i}\sqrt{1-\gamma},0,0) \tag{11}$$

$$\mathcal{T} = \mathrm{diag}(\sqrt{\gamma},\sqrt{\gamma},1,1) \tag{12}$$

We generate $\sim 10^6$ matrices $\mathcal{S}_0$ using the QR-based algorithm described in [22], and we build the corresponding matrices $\mathcal{S}$. Having extracted the submatrix $\mathbf{t}$ and diagonalized $\mathbf{t}\mathbf{t}^\dagger$, we collect its eigenvalues τ_1 and τ_2 and from them we construct the concurrence $\mathfrak{C}$. The resulting histogram is plotted in Fig. 2 for different values of γ together with the analytical result (8). The agreement is excellent.

In summary, we have investigated analytically the full distribution of concurrence for the entanglement production in a chaotic quantum dot supporting one ideal and one non-ideal lead. Our calculation is the first analytical result for the concurrence in non-ideal cavities. This result is in perfect agreement with the numerical findings in [15] (Fig. 6 bottom) and recovers, in the proper limit, the known results on the ideal transparency case [12]. In particular, we can confirm that the entanglement production is enhanced when the transparency of one of the contacts is far from ideal, and we have obtained also the analytical expression of the first moments, which may be tested experimentally. Moreover, we find that the structure of the probability distribution as a function of the transparency is simple and elegant, hinting towards a deeper connection with physical symmetries of the system. The comprehension of the generality of this relation when, for instance, both leads are non-ideal, is still lacking and represents a challenging issue whose solution is much called for.

Acknowledgments: The authors acknowledge the warm hospitality at the conference 'Non-equilibrium fluctuation-response relations' at Giglio Island, where this work was initiated. They also thank Carlo Barbieri of Wolfram|Alpha for useful discussions on some Mathematica routines.



FIG. 2: Probability density of the concurrence (Eq. (8)) compared with numerical simulations (black triangles) for different values of γ.



FIG. 3: Analytical predictions for the average and the standard deviation of concurrence as a function of γ (straight lines) together with numerical simulations (red circles).

[1] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).

[2] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).

[3] G. Alber, T. Beth, P. Horodecki, R. Horodecki, M. Rötteler, H. Weinfurter, R. Werner, and A. Zeilinger, *Quantum Information* (Springer, Berlin, 2001), Vol. 173.

[4] C. W. J. Beenakker, in *Quantum Computers, Algorithms and Chaos*, Proceedings of the International School of Physics "Enrico Fermi," Varenna, 2005, (IOS Press, Amsterdam, 2006).

[5] G. Burkard, J. Phys.: Condens. Matter **19**, 233202 (2007).

[6] P. Recher, E. V. Sukhorukov, and D. Loss, Phys. Rev. B **63**, 165314 (2001); G. B. Lesovik, T. Martin, and G. Blatter, Eur. Phys. J. B **24**, 287 (2001); W. D. Oliver, F. Yamaguchi, and Y. Yamamoto, Phys. Rev. Lett. **88**, 037901 (2002); V. Bouchiat, N. Chtchelkatchev, D. Feinberg, G. B. Lesovik, T. Martin, and J. Torres, Nanotechnology **14**, 77 (2003); D. S. Saraga and D. Loss, Phys. Rev. Lett. **90**, 166803 (2003).

[7] S. Bose and D. Home, Phys. Rev. Lett. **88**, 050401 (2002); C. W. J. Beenakker, C. Emary, M. Kindermann, and J. L. van Velsen, Phys. Rev. Lett. **91**, 147901 (2003); P. Samuelsson, E. V. Sukhorukov, and M. Büttiker, Phys. Rev. Lett. **92**, 026805 (2004) and New. J. Phys. **7**, 176 (2005); A. I. Signal and U. Zülicke, Appl. Phys. Lett. **87**, 102102 (2005); A. V. Lebedev, G. B. Lesovik, and G. Blatter, Phys. Rev. B **71**, 045306 (2005).

[8] D. Frustaglia, S. Montangero, and R. Fazio, Phys. Rev. B **74**, 165326 (2006).

[9] C. W. J. Beenakker, M. Kindermann, C. M. Marcus, and A. Yacoby, in *Fundamental Problems of Mesoscopic Physics*, edited by I. V. Lerner, B. L. Altshuler, and Y. Gefen, NATO Science Series II vol. 154 (Kluwer, Dordrecht, 2004).

[10] An orbital entangler for electrons based on an interacting mechanism was first introduced in P. Samuelsson, E. V. Sukhorukov, and M. Büttiker, Phys. Rev. Lett. **91**, 157002 (2003).

[11] F. Mezzadri and N. J. Simm, J. Math. Phys. **53**, 053504 (2012) and *ibid.* **52**, 103511 (2011); V. Al. Osipov and E. Kanzieper, Phys. Rev. Lett. **101**, 176804 (2008); B. A. Khoruzhenko, D. V. Savin, and H.-J. Sommers, Phys. Rev. B **80**, 125301 (2009); H.-J. Sommers, W. Wieczorek, and D. V. Savin, Acta Phys. Polon. A **112**, 691 (2007); M. Novaes, Phys. Rev. B **78**, 035337 (2008) and *ibid.* **75**, 073304 (2007); P. Vivo, S. N. Majumdar, and O. Bohigas, Phys. Rev. Lett. **101**, 216809 (2008) and Phys. Rev. B **81**, 104202 (2010); P. W. Brouwer and C. W. J. Beenakker, Phys. Rev. B **50**, 11263 (1994); C. W. J. Beenakker, Rev. Mod. Phys. **69**, 731 (1997); P. A. Mello and N. Kumar, *Quantum Transport in Mesoscopic Systems. Complexity and statistical fluctuations* (Oxford University Press, Oxford, 2004).

[12] V. A. Gopar and D. Frustaglia, Phys. Rev. B **77**, 153403 (2008).

[13] S. Rodriguez-Pérez and M. Novaes, Phys. Rev. B **85**, 205414 (2012).

[14] F. A. G. Almeida, S. Rodriguez-Perez, and A. M. S. Macêdo, Phys. Rev. B **80**, 125320 (2009).

[15] F. A. G. Almeida and A. M. C. Souza, Phys. Rev. B **82**, 115422 (2010).

[16] P. Vidal and E. Kanzieper, Phys. Rev. Lett. **108**, 206806 (2012).

[17] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).

[18] H. U. Baranger and P. A. Mello, Phys. Rev. Lett. **73**, 142 (1994).

[19] R. A. Jalabert, J.-L Pichard, and C. W. J. Beenakker, Europhys. Lett. **27**, 255 (1994).

[20] P. W. Brouwer, Phys. Rev. B **51**, 16878 (1995).

[21] See Supplemental Material.

[22] A. Edelman and N. Raj Rao, Acta Numerica **14**, 233 (2005).

Supplemental Material

We give here the form of the matrix $A_{ij}(\gamma)$ in the expansion of the joint probability density of eigenvalues $P_{\gamma,\gamma}(\tau_1,\tau_2)$ (Eq. (6) of the Letter).

$$A_{ij}(\gamma) = \begin{pmatrix} 0 & 0 & 6 & 12\gamma^2 & 2\gamma^4 \\ 0 & -12 & -12\gamma^2 & 52\gamma^4 & 12\gamma^6 \\ 6 & -12\gamma^2 & -108\gamma^4 & -12\gamma^6 & 6\gamma^8 \\ 12\gamma^2 & 52\gamma^4 & -12\gamma^6 & -12\gamma^8 & 0 \\ 2\gamma^4 & 12\gamma^6 & 6\gamma^8 & 0 & 0 \end{pmatrix} \tag{13}$$

We now start from Eq. (6) of the Letter for the joint probability density of transmission eigenvalues:

$$P_{\gamma,\gamma}(\tau_1,\tau_2) = \frac{(\gamma^2-1)^8 \sum_{i,j=0}^{4} A_{ij}(\gamma)(1-\tau_1)^i(1-\tau_2)^j}{\left(1-\gamma^2(1-\tau_1)\right)^6\left(1-\gamma^2(1-\tau_2)\right)^6} \tag{14}$$

and we have to compute the following integral:

$$P(\mathfrak{C}) = \int_{[0,1]^2} d\tau_1 d\tau_2 P_{\gamma,\gamma}(\tau_1,\tau_2)\delta\left(\mathfrak{C} - \frac{2\sqrt{\tau_1(1-\tau_1)\tau_2(1-\tau_2)}}{\tau_1+\tau_2-2\tau_1\tau_2}\right) \tag{15}$$

We make the change of variables (valid for $\gamma \neq 1$):

$$\begin{cases} (1-\gamma^2)z_1 &= \frac{\tau_1}{1-\tau_1} \\ (1-\gamma^2)z_2 &= \frac{\tau_2}{1-\tau_2} \end{cases} \tag{16}$$

This way, after simplifications, the integral becomes:

$$P(\mathfrak{C}) = \int_{[0,\infty]^2} dz_1 dz_2 \left[\sum_{k=0,2,4,6,8} \gamma^k f_k(z_1,z_2)\right] \delta\left(\mathfrak{C} - 2\frac{\sqrt{z_1 z_2}}{z_1+z_2}\right). \tag{17}$$

Note that the change of variable (16) is essential in bringing the polinomial structure in γ to the surface. The functions involved in Eq. (17) are:

$$f_0(z_1,z_2) = \frac{6(z_1-z_2)^2}{(1+z_1)^4(1+z_2)^4} \tag{18}$$

$$f_2(z_1,z_2) = -\frac{24(z_1-z_2)^2(-1+z_1z_2)}{(1+z_1)^5(1+z_2)^5} \tag{19}$$

$$f_4(z_1,z_2) = \frac{4(z_1-z_2)^2(15+z_1(6-4z_2)+6z_2-4z_2^2+z_1^2(-4+9z_2^2)}{(1+z_1)^6(1+z_2)^6} \tag{20}$$

$$f_6(z_1,z_2) = -\frac{8(z_1-z_2)^2(-3-(-6+z_2)z_2+z_1(6+(8-3z_2)z_2)+z_1^2(-1+3(-1+z_2)z_2)}{(1+z_1)^6(1+z_2)^6} \tag{21}$$

$$f_8(z_1,z_2) = \frac{2(z_1-z_2)^2(3+(-6+z_2)z_2+z_1^2(1+3(-2+z_2)z_2)+z_1(-6+28z_2-6z_2^2)}{(1+z_1)^6(1+z_2)^6} \tag{22}$$

The integrals of the form:

$$c_k(\mathfrak{C}) = \int_{[0,\infty]^2} dz_1 dz_2 f_k(z_1,z_2)\delta\left(\mathfrak{C} - 2\frac{\sqrt{z_1z_2}}{z_1+z_2}\right) \tag{23}$$

can be computed by first expanding the delta function as:

$$\delta\left(\mathfrak{C} - 2\frac{\sqrt{z_1z_2}}{z_1+z_2}\right) = \frac{(\phi^{(+)}(\mathfrak{C})+1)^2\sqrt{\phi^{(+)}(\mathfrak{C})}}{\phi^{(+)}(\mathfrak{C})-1} z_2\, \delta(z_1 - \phi^{(+)}(\mathfrak{C})z_2) + \frac{(\phi^{(-)}(\mathfrak{C})+1)^2\sqrt{\phi^{(-)}(\mathfrak{C})}}{1-\phi^{(-)}(\mathfrak{C})} z_2\, \delta(z_1 - \phi^{(-)}(\mathfrak{C})z_2) \tag{24}$$

where

$$\phi^{(\pm)}(\mathfrak{C}) = \frac{2-\mathfrak{C}^2 \pm 2\sqrt{1-\mathfrak{C}^2}}{\mathfrak{C}^2} \tag{25}$$

yielding eventually:

$$c_k(\mathfrak{C}) = \frac{(\phi^{(+)}(\mathfrak{C})+1)^2\sqrt{\phi^{(+)}(\mathfrak{C})}}{\phi^{(+)}(\mathfrak{C})-1} c_k^{(+)}(\mathfrak{C}) + \frac{(\phi^{(-)}(\mathfrak{C})+1)^2\sqrt{\phi^{(-)}(\mathfrak{C})}}{1-\phi^{(-)}(\mathfrak{C})} c_k^{(-)}(\mathfrak{C}) \tag{26}$$

where:

$$c_k^{(\pm)}(\mathfrak{C}) = \int_0^\infty dz\; z\; f_k\left(\phi^{(\pm)}(\mathfrak{C})z, z\right) \tag{27}$$

The single integrals in (27) can be computed and after lengthy algebra we eventually get to the following coefficients:

$$c_0(\mathfrak{C}) = \frac{\mathfrak{C}\left(4\sqrt{1-\mathfrak{C}^2}(11+4\,\mathfrak{C}^2) + (6+9\,\mathfrak{C}^2)\ln\left(\frac{2-\mathfrak{C}^2-2\sqrt{1-\mathfrak{C}^2}}{2-\mathfrak{C}^2+2\sqrt{1-\mathfrak{C}^2}}\right)\right)}{2(-1+\mathfrak{C}^2)^3} \tag{28}$$

$$\begin{aligned} c_4(\mathfrak{C}) = \frac{\mathfrak{C}}{3(\mathfrak{C}^2-1)^4}&\left(4\sqrt{1-\mathfrak{C}^2}(62+221\,\mathfrak{C}^2+32\,\mathfrak{C}^4) + 6(8+64\,\mathfrak{C}^2+33\,\mathfrak{C}^4)\times\right. \\ &\left.\times\text{arctanh}\left(\frac{2\sqrt{1-\mathfrak{C}^2}}{\mathfrak{C}^2-2}\right)\right) \end{aligned} \tag{29}$$

$$\begin{aligned} c_6(\mathfrak{C}) = -\frac{\mathfrak{C}}{6(\mathfrak{C}^2-1)^4}&\left(4\sqrt{1-\mathfrak{C}^2}(62+221\mathfrak{C}^2+32\,\mathfrak{C}^4)+\right. \\ &\left.3(8+64\,\mathfrak{C}^2+33\,\mathfrak{C}^4)\ln\left(\frac{2-\mathfrak{C}^2-2\sqrt{1-\mathfrak{C}^2}}{2-\mathfrak{C}^2+2\sqrt{1-\mathfrak{C}^2}}\right)\right) \end{aligned} \tag{30}$$

$$\begin{aligned} c_8(\mathfrak{C}) = \frac{\mathfrak{C}}{12(\mathfrak{C}^2-1)^5}&\left(2\sqrt{1-\mathfrak{C}^2}(80+1212\,\mathfrak{C}^2+1431\,\mathfrak{C}^4+112\,\mathfrak{C}^6)+\right. \\ &\left.3(8+236\,\mathfrak{C}^2+554\,\mathfrak{C}^4+147\,\mathfrak{C}^6)\text{arctanh}\left(\frac{2\sqrt{1-\mathfrak{C}^2}}{\mathfrak{C}^2-2}\right)\right) \end{aligned} \tag{31}$$

Note that the coefficient $c_2(\mathfrak{C})$ is identically zero.
Now, consider the general expansion of the joint probability distribution function in Eq. (6) of the Letter

$$P_{\gamma,\gamma}(\tau_1,\tau_2) = \sum_{k=0}^{+\infty} \gamma^{2k} P^{(2k)}(\tau_1,\tau_2). \tag{32}$$

where in particular

$$P^{(0)}(\tau_1,\tau_2) = 48(\tau_1-\tau_2)^2(1-(\tau_1+\tau_2)). \tag{33}$$

Note that under the transformations

$$\begin{cases} \tau_1 & \to 1-\tau_1 \\ \tau_2 & \to 1-\tau_2 \end{cases} \tag{34}$$

$P^{(0)}(\tau_1,\tau_2)$ is antisymmetric and therefore it is straightforward to observe that:

$$\langle \mathfrak{C}^n \rangle = \int d\tau_1 d\tau_2 P^{(0)}(\tau_1,\tau_2)\left[\frac{2\sqrt{\tau_1(1-\tau_1)\tau_2(1-\tau_2)}}{\tau_1+\tau_2-2\tau_1\tau_2}\right]^n \equiv 0 \tag{35}$$

for all values of n. Note that also for higher order terms than $\mathcal{O}(\gamma^8)$, all the integrals of the kind:

$$\langle \mathfrak{C}^n \rangle = \int d\tau_1 d\tau_2 P^{(2k)}(\tau_1, \tau_2) \left[\frac{2\sqrt{\tau_1(1-\tau_1)\tau_2(1-\tau_2)}}{\tau_1 + \tau_2 - 2\tau_1\tau_2} \right]^n \tag{36}$$

vanish for all $k > 4$ and every n , as can be clearly deduced from the exact probability density of the concurrence $\mathfrak{C}$. However, quite interestingly, the simple symmetry argument (34) or other simple transformations do not hold for these terms.

In conclusion, moments of arbitrary order can be computed exactly, also from the expansion (32), if needed. For instance, the first two are:

$$\overline{\mathfrak{C}} = a_0 + a_4\gamma^4 + a_6\gamma^6 + a_8\gamma^8 \tag{37}$$

$$\overline{\mathfrak{C}^2} = b_0 + b_4\gamma^4 + b_6\gamma^6 + b_8\gamma^8 \tag{38}$$

$$a_0 = \frac{1}{16}\pi(21\pi - 64) \tag{39}$$

$$a_4 = \frac{1}{48}\pi(327\pi - 1024) \tag{40}$$

$$a_6 = -\frac{1}{96}\pi(327\pi - 1024) \tag{41}$$

$$a_8 = \frac{\pi(9129\pi - 28672)}{3072} \tag{42}$$

$$b_0 = -22 + \frac{9\pi^2}{4} \tag{43}$$

$$b_4 = \frac{1}{6}\left(-976 + 99\pi^2\right) \tag{44}$$

$$b_6 = \frac{244}{3} - \frac{33\pi^2}{4} \tag{45}$$

$$b_8 = \frac{147\pi^2}{16} - \frac{272}{3} \tag{46}$$